

Product	Dosage instructions
AIR REPAIR 250ml	Spray Air Repair into the air or onto contaminated areas as necessary
ARTHRIMED TABLETS 30's, 90's	<p>Dosage - dogs:</p> <p>Small dogs: (1-10kg) 1 tablet per day Medium dogs: (11-20kg) 2 tablets per day Large dogs: (21-40kg) 3 tablets per day Giant dogs: (41+ kg) 4 tablets per day</p> <p>Dosage - cats:</p> <p>Small cats: (1-3kg) ½ tablet per day Large cats: (4+ kg) 1 tablet per day</p>
ARTHRIMED POWDER 250g	<p>Dosage - dogs:</p> <p>Small dogs: (1-10kg) 5g per day Medium dogs: (11-20kg) 2 x 5g per day Large dogs: (21-40kg) 3 x 5g per day Giant dogs: (41+ kg) 4 x5g per day</p> <p>Dosage - cats:</p> <p>Small cats: (1-3kg) 2,5g per day Large cats: (4+ kg) 5g per day</p>
ARTHRITIS CARE TABLETS 60's	<p>Dosage - dogs:</p> <p>Small dogs: (1-10kg) 1 tablet per day Medium dogs: (11-20kg) 2 tablets per day Large dogs: (21-40kg) 3 tablets per day Giant dogs: (41+ kg) 4 tablets per day</p> <p>Dosage - cats:</p> <p>Small cats: (1-3kg) ½ tablet per day Large cats: (4+ kg) 1 tablet per day</p>
BIO-VITA 200g, 400g, 5kg	<p>Dogs: 5g (1 measure) per 10kg of bodyweight per day. Double the dosage for pregnant and lactating bitches.</p> <p>Puppies: ½ measure (2,5g) per day</p> <p>Cats: 5g (1 measure) per day.</p> <p>Kittens: ½ measure (2,5g) per day</p>
CA-DOG 250g, 500g	<p>Sprinkle one to two 5g measures on the food daily.</p> <p>Increase the dose for pregnant or lactating bitches.</p> <p>This product is not recommended for cats.</p>
DICESTAL TABS 10's, 100's	<p>By oral administration preferably just before the main meal of the day.</p> <p>1 x 0,5 g tablet per 2,5 kg body mass</p>
DOXYDOG 100'S (Y) 50mg	Dose 1 x 50 mg DoxyDog (yellow tablet) per 5 kg bodymass daily for 10 - 20 days. (Do not break the tablet).
DOXYDOG 100' S (R) 100mg	Dose 1 x 100 mg DoxyDog (red tablet) per 10 kg bodymass once daily for 10 – 20 days. (Do not break the tablet).
DOXYDOG 50' S (O) 300mg	Dose 1 x 300 mg Doxydog (orange tablet) per 30 kg bodymass once daily for 10 – 20 days. (Do not break the tablet).
DOXYSYRUP 25ml	Dose 1 ml of Doxysyrup per kg body mass once daily for 5 to 7 days
ESBILAC MILK REPLACER 5 PDS **	<p>Gently shake or stir one part of powdered Esbilac into two parts of warm water. (A part is any measuring device you use: teaspoon, tablespoon, or cup.)</p> <p>Do not mix more Esbilac than can be consumed in 24 hours.</p> <p>Do not use a blender.</p> <p>Reconstituted Esbilac should be kept refrigerated.</p>
FLY REPEL 250ml	<p>WIPE - ON USE:</p> <p>Brush animal's coat to remove excess dirt and dust. Shake container well and dampen a sponge or soft cloth with Fly Repel.</p> <p>Rub over dog's hair with special attention to legs, shoulders, neck and facial area, avoiding eyes.</p> <p>Dab, smear or paint lightly onto dog's ears and other areas where protection is required. Repeat as necessary.</p> <p>SPRAY – ON USE:</p> <p>As above, but spray onto animal instead of wiping on. Do not spray mist around the head and eyes. Dampen a soft cloth and wipe head avoiding eyes.</p> <p>For treatment of ticks - Fly Repel can be used as a spot treatment spray, or applied as described above for total body protection.</p>
HEPAVET TABLETS 100's	<p>Small dogs: 1 tablet daily</p> <p>Medium dogs: 2 tablets daily</p>

	Large dogs: 4 tablets daily
HYGIENE SHAMPOO	Wet coat. Apply Medpet Hygiene Shampoo onto coat and work to lather. Rinse well with water.
INTESTIPET	Add 10g (2 heaped measures) to pet's food, per 2.25kg live weight. Moisten food with water or edible oil to allow Intestipet to adhere to the food Routine treatment: Administer routinely once a week During Antibiotic treatment: Administer daily over the food during the antibiotic treatment. During Gastro-intestinal disease: Administer daily while symptoms persist. Administer for 2 additional days.
MEDICOTE 100ml, refill	DOGS: Toy breeds: (1-5kg) 1 pump stroke Small dogs: (6-10kg) 2 pump strokes Medium dogs: (11-20kg) 3 pump strokes Large dogs: (21-40kg) 4 pump strokes Giant dogs: (41+ kg) 5 pump strokes CATS: Small cats: (1-3kg) 1 pump stroke Large cats: (4+ kg) 2 pump strokes HORSES: Medium breeds: (up to 300kg) 10 pump strokes Large breeds: (300+ kg) 15 pump strokes
MEDI-GUT 100ML	Shake well before use. Initial dose: 4 squirts over food or in mouth three times a day Maintenance dose: 4 squirts over food or in mouth once a day.
MEDIMUNE TABLETS 30's	Dogs: 1 tablets per 10kg Small cats: (1-3kg) ½ tablet Large cats: (4+ kg) 1 tablet
MEDIWORM PLUS 100's	1 tablet per 10 kg body mass
MEDIWORM FOR CATS 25's	1 tablet per 5kg bodymass
MEDIWORM FOR DOGS 100's	1 tablet per 10 kg body mass
MEDIWORM FOR LARGE DOGS 15's	1 tablet per 40 kg bodymass
OROVET ORAL SPRAY/RINSE 250ml	Apply 10ml 2x daily to the gum (gingival) and inside (mucosa) of the lips for at least 10days. Do not rinse the animal's mouth after applications or give it water for at least 10-20 minutes. Do not apply gel to the animal's mouth 1 hour before or after a meal.
OROZYME GEL TUBE 70g	Use 1 or 2 times per week (brush the pet's teeth if possible).
PET ASSIST 1kg	Administer 2g (½ scoop per 5kg of bodymass
RIGLY FART-EZE 250g	Add ½ level measuring spoon (2,5g) per 10kg bodymass with the dog's food at each meal
RX FOR FLEAS 400g	Apply directly on carpets wherever pets are frequently found or sleep at a rate of 500g per 5 square meters. Work the powder deep into the fibres and the mat allowing it to remain for a period of three weeks to achieve maximum flea control. No powder should be visible after application. Powder that does not brush in completely must be vacuumed immediately.
SALMON OIL 100ml	Dogs: 1ml per 5kg bodyweight Cats: 1ml per 3kg bodyweight Horses: 1ml per 30kg bodyweight
TEA TREE OIL SHAMPOO 250ml	Wet coat thoroughly. Apply shampoo liberally and work into a rich lather. Rinse thoroughly.
VITA-KELP 250g, 500g	1 teaspoon = 5 g. Sprinkle on the food daily. Kittens: 1,25g (¼ teaspoonfull) daily Cats, puppies, small dogs: 2,5g (½ teaspoonfull) daily Medium dogs: 5g (1 teaspoonfull) daily Large dogs: 10g (2 teaspoonsfull) daily
M & B TABLETS 50's, 250's	2 Tablets per 5 kg body mass followed by half initial dose daily for a maximum of four days.
PHENAMIDINE INJ 25ml	Administer 0.3ml per 1kg body mass, by subcutaneous injection.